Author index

Abe, A. and Sasaki, T.
Sulfhydryl groups in glycolipid transfer protein: formation of an intramolecular disulfide bond and oligomers by Cu^{2+}-catalyzed oxidation (985) 38
Abe, A. and Sasaki, T.
Formation of an intramolecular disulfide bond of glycolipid transfer protein (985) 45
Akanuma, H., see Sakai, K. (985) 51
Aranda, F.J., Coutinho, A., Berberan-Santos, M.N., Prieto, M.J.E. and Gómez-Fernández, J.C.
Fluorescence study of the location and dynamics of α-tocopherol in phospholipid vesicles (985) 26
Artigues, A., Villar, M.T., Fernandez, A.M., Ferragut, J.A. and Gonzalez-Ros, J.M.
Cholesterol stabilizes the structure of the nicotinic acetylcholine receptor reconstituted in lipid vesicles (985) 325
Atkins, A.R., see Cornell, B.A. (985) 229

Barenholz, Y., see Ben-Yashar, V. (985) 271
Bartel, D., Hans, H. and Passow, H.
Identification by site-directed mutagenesis of Lys-558 as the covalent attachment site of H_2DIDS in the mouse erythroid band 3 protein (985) 355
Basu, J., see Kundu, M. (985) 97
Ben-Yashar, V. and Barenholz, Y.
The interaction of cholesterol and cholest-4-en-3-one with dipalmitoylphosphatidylcholine. Comparison based on the use of three fluorophores (985) 271
Benkouka, F., see Hervatin, F. (985) 320
Berberan-Santos, M.N., see Aranda, F.J. (985) 26
Blank, M., see Korbmacher, C. (985) 67
Boggs, J.M., Wang, H.-Y., Rangaraj, G. and Tümmler, B.
Interdigitation of phosphatidylcholine and phosphatidylethanolamine mixed with complexes of acidic lipids and polymyxin B or polymyxin B nonapeptide (985) 199
Boggs, J.M., see Wang, H.-Y. (985) 182
Bourre, J.-M., see Tayarani, I. (985) 168
Brajtburg, J., Gruda, I., Daigle, I. and Medoff, G.
Concentration dependent dual effect of the monolauryl ester of sucrose on the antifungal activity and absorption spectra of amphotericin B (Fungizone) (985) 307

Carruthers, A., see Naderi, S. (985) 173
Chakrabarti, P., see Kundu, M. (985) 97
Chung, Y.J. and Jeffrey, G.A.
The lyotropic liquid crystal properties of *n*-octyl 1-*O*-β-D-glucopyranoside and related *n*-alkyl pyranosides (985) 300
Cloëz, I., see Tayarani, I. (985) 168
Cornell, B.A., Separovic, F., Thomas, D.E., Atkins, A.R. and Smith, R.
Effect of acyl chain length on the structure and motion of gramicidin A in lipid bilayers (985) 229
Coutinho, A., see Aranda, F.J. (985) 26

Daigle, I., see Brajtburg, J. (985) 307
De Cock, H., Meeldijk, J., Overduin, P., Verkleij, A. and Tommassen, J.
Membrane biogenesis in *Escherichia coli*: effects of a *secA* mutation (985) 313
Donath, E., see Engel, J. (985) 111
Dressler, S., see Korbmacher, C. (985) 67
Duerre, J.A. and Gordon, R.K.
Characterization of the binding of *S*-adenosyl-L-methionine to plasma membranes of HL-60 promyelocytic leukemia cells (985) 211

Eisenthal, R., Game, S. and Holman, G.D.
Specificity and kinetics of hexose transport in *Trypanosoma brucei* (985) 81
Ekman, S., see Slotte, J.P. (985) 90
Engel, J., Donath, E., Ermakov, Y.A., Meyer, H.W. and Richter, W.
Hyperosmotic relaxation lysis of chromaffin granules is caused by interactions between the granular membrane and intragranular vesicles (985) 111
Ermakov, Y.A., see Engel, J. (985) 111

Fernandez, A.M., see Artigues, A. (985) 325
Fernandez-Belda, F., Soler, F. and Gomez-Fernandez, J.C.
Quinacrine inhibits the calcium-induced calcium release in heavy sarcoplasmic reticulum vesicles (985) 279
Ferragut, J.A., see Artigues, A. (985) 325
Fischer, T.M.
Cross bonding and stiffening of the red cell membrane (985) 218

Game, S., see Eisenthal, R. (985) 81
Ganapathy, V., see Kulanthaivel, P. (985) 139
Glaser-Wuttke, G., Keppner, J. and Rasched, I.
Pore-forming properties of the adsorption protein of filamentous phage fd (985) 239
Gómez-Fernández, J.C., see Aranda, F.J. (985) 26
Gomez-Fernandez, J.C., see Fernandez-Belda, F. (985) 279
Gonzalez-Ros, J.M., see Artigues, A. (985) 325
Gordon, R.K., see Duerre, J.A. (985) 211
Grell, E., see Kojro, Z. (985) 1
Gruda, I., see Brajtburg, J. (985) 307
Guerin, M., see Manon, S. (985) 127

Hamada, A., see Naito, M. (985) 147
Hans, H., see Bartel, D. (985) 355
Hayashi, M., see Sakai, K. (985) 51
Hedström, G., see Slotte, J.P. (985) 90
Helbig, H., see Korbmacher, C. (985) 67
Hervatin, F., Benkouka, F., Robert, J.C., Péranzi, G. and Soumarmon, A.
The ontogeny of rat gastric H^+/K^+-ATPase (985) 320
Hicks, D.B., see Rohde, M. (985) 233
Higuchi, A., see Kominami, S. (985) 293
Holman, G.D., see Eisenthal, R. (985) 81
Huang, L., see Pinnaduwage, P. (985) 33
Hunt, G.R.A., see Kaszuba, M. (985) 106

Inoue, K., see Naito, M. (985) 147
Inoue, S., see Kominami, S. (985) 293

Jansen, G., Westerhof, G.R., Rijksen, G. and Schornagel, J.H.
Interaction of *N*-hydroxy(sulfo)succinimide active esters with the reduced folate/methotrexate transport system from human leukemic CCRF-CEM cells (985) 266

Jeffrey, G.A., see Chung, Y.J. (985) 300
Jensen, J. and Nørby, J.G.
Thallium binding to native and radiation-inactivated Na^+/K^+-ATPase (985) 248

Karlish, S.J.D., see Tal, D.M. (985) 55
Kaszuba, M. and Hunt, G.R.A.
A ^{1}H-NMR study of the influence of *n*-alcohols on the stoichiometry of melittin-induced permeability of phospholipid vesicles (985) 106
Kates, M., see Pugh, E.L. (985) 255
Kawamura, Y., see Sakurai, I. (985) 347
Kawasaki, T., see Satoh, O. (985) 120
Kawashima, S., see Sakai, K. (985) 51
Keppner, J., see Glaser-Wuttke, G. (985) 239
Knight, C.G. and Matthews, A.
Determination of pK_a values of lipophilic fluorescein derivatives in phospholipid vesicles by fluorescence emission ratio titration (985) 75
Koepsell, H., see Rausch, T. (985) 133
Koike, H., see Kora, S. (985) 286
Kojro, Z., Lin, S.-Q., Grell, E. and Ruf, H.
Determination of internal volume and volume distribution of lipid vesicles from dynamic light scattering data (985) 1
Kominami, S., Inoue, S., Higuchi, A. and Takemori, S.
Steroidogenesis in liposomal system containing adrenal microsomal cytochrome *P*-450 electron transfer components (985) 293
Kora, S., Sado, M., Koike, H. and Terada, H.
Protective effect of the plasticizer di(2-ethylhexyl) phthalate against damage of the mitochondrial membrane induced by calcium: possible participation of the adenine nucleotide translocator (985) 286
Korbmacher, C., Helbig, H., Blank, M., Dressler, S., Stahl, F. and Wiederholt, M.
Characterization of Cl^-/HCO_3^- exchange in A10 vascular smooth muscle cells using $^{36}Cl^-$ (985) 67
Krajcarski, D.T., see Pugh, E.L. (985) 255
Krulwich, T.A., see Rohde, M. (985) 233
Kudo, I., see Naito, M. (985) 147
Kudo, Y., see Satoh, O. (985) 120
Kulanthaivel, P., Leibach, F.H., Mahesh, V.B. and Ganapathy, V.
Tyrosine residues are essential for the activity of the human placental taurine transporter (985) 139
Kundu, M., Basu, J., Rakshit, M.M. and Chakrabarti, P.
Abnormalities in erythrocyte membrane band 3 in chronic myelogenous leukemia (985) 97

Lakey, J.H., Maget-Dana, R. and Ptak, M.
The lipopeptide antibiotic A21978C has a specific interaction with DMPC only in the presence of calcium ions (985) 60
Lefauconnier, J.-M., see Tayarani, I. (985) 168
Leibach, F.H., see Kulanthaivel, P. (985) 139
Lin, S.-Q., see Kojro, Z. (985) 1
Lönnendonker, U.
Use-dependent block of sodium channels in frog myelinated nerve by tetrodotoxin and saxitoxin at negative holding potentials (985) 153
Lönnendonker, U.
Binding of tetrodotoxin and saxitoxin to Na^+ channels at different holding potentials: fluctuation measurements in frog myelinated nerve (985) 161

Maget-Dana, R., see Lakey, J.H. (985) 60
Mahesh, V.B., see Kulanthaivel, P. (985) 139
Manon, S. and Guerin, M.
Modifications of the relative proteolipid composition in the ATP synthase of a respiratory competent mutant of *Saccharomyces cerevisiae* (985) 127
Matthews, A., see Knight, C.G. (985) 75
Mayer, F., see Rohde, M. (985) 233
Medoff, G., see Brajtburg, J. (985) 307
Meeldijk, J., see De Cock, H. (985) 313
Melchior, D.L., see Naderi, S. (985) 173
Menestrina, G., see Ropele, M. (985) 9
Meyer, H.W., see Engel, J. (985) 111
Mishima, K. and Morimoto, T.
Electric field-induced orientation of myelin figures of phosphatidylcholine (985) 351
Morimoto, T., see Mishima, K. (985) 351
Morjana, N.A. and Scarborough, G.A.
Evidence for an essential histidine residue in the *Neurospora crassa* plasma membrane H^+-ATPase (985) 19

Naderi, S., Carruthers, A. and Melchior, D.L.
Modulation of red blood cell sugar transport by lyso-lipid (985) 173
Naito, M., Utsumi, H., Umeda, M., Kudo, I., Takeshita, K., Hamada, A., Nojima, S. and Inoue, K.
ESR study on synthetic glyceroglycolipid liposomal membranes (985) 147
Nojima, S., see Naito, M. (985) 147
Nørby, J.G., see Jensen, J. (985) 248

Overduin, P., see De Cock, H. (985) 313

Passow, H., see Bartel, D. (985) 355
Péranzi, G., see Hervatin, F. (985) 320
Pinnaduwage, P., Schmitt, L. and Huang, L.
Use of a quaternary ammonium detergent in liposome mediated DNA transfection of mouse L-cells (985) 33
Prieto, M.J.E., see Aranda, F.J. (985) 26
Ptak, M., see Lakey, J.H. (985) 60
Pugh, E.L., Kates, M., Szabo, A.G. and Krajcarski, D.T.
Correlation of enzyme activities with fluorescence anisotropy of dansyl-labeled cytochrome b_5/NADH–cytochrome-b_5 reductase systems in phosphatidylcholine vesicles (985) 255

Rakshit, M.M., see Kundu, M. (985) 97
Rangaraj, G., see Boggs, J.M. (985) 199
Rannström, S., see Slotte, J.P. (985) 90
Rasched, I., see Glaser-Wuttke, G. (985) 239
Raszeja-Specht, A., see Rausch, T. (985) 133
Rausch, T., Raszeja-Specht, A. and Koepsell, H.
Identification of an M_r 75000 component of the H^+/D-glucose cotransporter from *Zea mays* with monoclonal antibodies directed against the mammalian Na^+/D-glucose cotransporter (985) 133
Richter, W., see Engel, J. (985) 111
Rijksen, G., see Jansen, G. (985) 266
Robert, J.C., see Hervatin, F. (985) 320
Rohde, M., Mayer, F., Hicks, D.B. and Krulwich, T.A.
Immunoelectron microscopic localization of the F_1F_0 ATPase (ATP synthase) on the cytoplasmic membrane of alkalophilic *Bacillus firmus* RAB (985) 233
Ropele, M. and Menestrina, G.
Electrical properties and molecular architecture of the channel formed by *Escherichia coli* hemolysin in planar lipid membranes (985) 9
Ruf, H., see Kojro, Z. (985) 1

Sado, M., see Kora, S. (985) 286
Sakai, K., Hayashi, M., Kawashima, S. and Akanuma, H.
Calcium-induced localization of calcium-activated neutral proteinase on plasma membranes (985) 51
Sakurai, I. and Kawamura, Y.
Effect of a magnetic field on charge flow along the interface between phosphatidylcholine monolayer and water (985) 347

Sakurai, I., Suzuki, T. and Sakurai, S.
Cross-sectional view of myelin figures (985) 101
Sakurai, S., see Sakurai, I. (985) 101
Sasaki, T., see Abe, A. (985) 38
Sasaki, T., see Abe, A. (985) 45
Satoh, O., Kudo, Y., Shikata, H., Yamada, K. and Kawasaki, T.
Characterization of amino-acid transport systems in guinea-pig intestinal brush-border membrane (985) 120
Scarborough, G.A., see Morjana, N.A. (985) 19
Schmitt, L., see Pinnaduwage, P. (985) 33
Schornagel, J.H., see Jansen, G. (985) 266
Separovic, F., see Cornell, B.A. (985) 229
Shikata, H., see Satoh, O. (985) 120
Slotte, J.P., Hedström, G., Rannström, S. and Ekman, S.
Effects of sphingomyelin degradation on cell cholesterol oxidizability and steady-state distribution between the cell surface and the cell interior (985) 90
Smith, R., see Cornell, B.A. (985) 229
Soler, F., see Fernandez-Belda, F. (985) 279
Soumarmon, A., see Hervatin, F. (985) 320
Sowers, A.E.
Evidence that electrofusion yield is controlled by biologically relevant membrane factors (985) 334
Sowers, A.E.
Electrofusion of dissimilar membrane fusion partners depends on additive contributions from each of the two different membranes (985) 339
Stahl, F., see Korbmacher, C. (985) 67
Suzuki, T., see Sakurai, I. (985) 101
Szabo, A.G., see Pugh, E.L. (985) 255

Takahashi, K., see Yoshida, T. (985) 331
Takemori, S., see Kominami, S. (985) 293
Takeshita, K., see Naito, M. (985) 147
Tal, D.M., Yanuck, M.D., Van Hall, G. and Karlish, S.J.D.
Identification of Na^+/K^+-ATPase inhibitors in bovine plasma as fatty acids and hydrocarbons (985) 55
Tayarani, I., Cloëz, I., Lefauconnier, J.-M. and Bourre, J.-M.
Sodium-dependent high-affinity uptake of taurine by isolated rat brain capillaries (985) 168
Terada, H., see Kora, S. (985) 286
Thomas, D.E., see Cornell, B.A. (985) 229
Tommassen, J., see De Cock, H. (985) 313
Tse, C.-M. and Young, J.D.
Passive and carrier-mediated permeation of different nucleosides through the reconstituted nucleoside transporter (985) 343
Tümmler, B., see Boggs, J.M. (985) 199
Tümmler, B., see Wang, H.-Y. (985) 182

Ueda, I., see Yoshida, T. (985) 331
Umeda, M., see Naito, M. (985) 147
Utsumi, H., see Naito, M. (985) 147

Van Hall, G., see Tal, D.M. (985) 55
Verkleij, A., see De Cock, H. (985) 313
Villar, M.T., see Artigues, A. (985) 325

Wang, H.-Y., Tümmler, B. and Boggs, J.M.
Use of spin labels to determine the percentage of interdigitated lipid in complexes with polymyxin B and polymyxin B nonapeptide (985) 182
Wang, H.-Y., see Boggs, J.M. (985) 199
Westerhof, G.R., see Jansen, G. (985) 266
Wiederholt, M., see Korbmacher, C. (985) 67

Yamada, K., see Satoh, O. (985) 120
Yanuck, M.D., see Tal, D.M. (985) 55
Yoshida, T., Takahashi, K. and Ueda, I.
Molecular orientation of volatile anesthetics at the binding surface: ^1H- and ^{19}F-NMR studies of submolecular affinity (985) 331
Young, J.D., see Tse, C.-M. (985) 343